

COMBINATORIAL AND MODEL-THEORETICAL PRINCIPLES RELATED TO REGULARITY OF ULTRAFILTERS AND COMPACTNESS OF TOPOLOGICAL SPACES. V.

PAOLO LIPPARINI

ABSTRACT. We generalize to the relations $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ and $\text{alm}(\lambda, \mu) \xrightarrow{\kappa} \text{alm}(\lambda', \mu')$ some results obtained in Parts II and IV. We also present a multi-cardinal version.

In this note we present a version of [L5, II, Theorem 1] and [L5, IV, Theorem 7] for the relation $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ and some variants.

See Parts I, II, IV [L5] or [BF, CN, CK, KM, L1, L2, L3] for unexplained notation.

1. EQUIVALENTS OF $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$

Let us recall the definition of $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$, as given in [L4]. Another formulation (equivalent for $\kappa \geq \sup\{\lambda, \lambda'\}$) had been given in [L1]. Notice that the two formulations are **not necessarily equivalent** for $\kappa < \sup\{\lambda, \lambda'\}$. See also [L3, Section 0], and Definition 1.2 and Theorem 1.3 below.

Given infinite cardinals $\lambda \geq \mu$ and a set $y \in S_\mu(\lambda)$, we denote by $[y]_{S_\mu(\lambda)}$, or simply by $[y]$ if there is no danger of confusion, the “cone” $\{s \in S_\mu(\lambda) \mid y \subseteq s\}$ of y in $S_\mu(\lambda)$.

Definition 1.1. [L4] We say that an ultrafilter D over $S_\mu(\lambda)$ *covers* λ if and only if $[\{\alpha\}] = \{s \in S_\mu(\lambda) \mid \alpha \in s\} \in D$, for every $\alpha \in \lambda$.

Let $\lambda \geq \mu$, $\lambda' \geq \mu'$ be infinite cardinals, and κ be any cardinal.

The relation $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ holds if and only if there are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that whenever D is an ultrafilter over

2000 *Mathematics Subject Classification.* Primary 03C20, 03E05, 03C95; Secondary 03C55, 03C98.

Key words and phrases. Regular, almost regular elementary extensions of models with power sets as a base; infinite matrices of sets; regular ultrafilters; images of ultrafilters; compact abstract logics.

The author has received support from MPI and GNSAGA. We wish to express our gratitude to X. Caicedo for stimulating discussions and correspondence.

$S_\mu(\lambda)$, and D covers λ , then for some $\beta \in \kappa$ it happens that $f_\beta(D)$ covers λ' .

In order to state the next theorem, we must recall some definitions from [L1] and [L3, Section 0].

Definition 1.2. $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ denotes the model $\langle S_{\mu''}(\lambda''), \subseteq, U, U', \{\alpha\} \rangle_{\alpha \in \lambda''}$, where $\lambda'' = \sup\{\lambda, \lambda'\}$, $\mu'' = \sup\{\mu, \mu'\}$, $U(x)$ if and only if $x \in S_\lambda(\mu)$, and $U'(y)$ if and only if $y \in S_{\lambda'}(\mu')$.

If $\mathfrak{S}^+$ is an expansion of $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$, and $\mathfrak{B} \equiv \mathfrak{S}^+$, we say that $\mathfrak{B}$ is (λ, μ) -regular if and only if there is $b \in B$ such that $\mathfrak{B} \models U(b)$, and $\mathfrak{B} \models \{\alpha\} \subseteq b$ for every $\alpha \in \lambda$.

Similarly, $\mathfrak{B}$ is (λ', μ') -regular if and only if there is $b' \in B$ such that $\mathfrak{B} \models U'(b')$, and $\mathfrak{B} \models \{\alpha'\} \subseteq b'$ for every $\alpha' \in \lambda'$.

Theorem 1.3. Suppose that $\lambda \geq \mu$, $\lambda' \geq \mu'$ are infinite cardinals, and κ is any cardinal.

Then the following conditions are equivalent.

- (a) $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ holds.
- (b) There are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that for every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \subseteq \bigcup_{\beta \in F} f_\beta^{-1}[\{g(\beta)\}]$.
- (c) There is a family $(C_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:
 - (i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcap_{\alpha \in H} C_{\alpha, \beta} = \emptyset$.
 - (ii) For every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \subseteq \bigcup_{\beta \in F} C_{g(\beta), \beta}$.
- (c') There is a family $(B_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:
 - (i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcup_{\alpha \in H} B_{\alpha, \beta} = S_\mu(\lambda)$.
 - (ii) For every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \cap \bigcap_{\beta \in F} B_{g(\beta), \beta} = \emptyset$.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, then the preceding conditions are also equivalent to:

(d) $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ has a multi-sorted expansion $\mathfrak{S}^+$ with at most κ new symbols such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is (λ, μ) -regular, then $\mathfrak{B}$ is (λ', μ') -regular.

(e) $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ has an expansion $\mathfrak{S}^+$ with at most κ new symbols such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is (λ, μ) -regular, then $\mathfrak{B}$ is (λ', μ') -regular.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, and $\lambda' = \mu'$ is a regular cardinal, then the preceding conditions are also equivalent to:

(f) Every κ -(λ, μ)-compact logic is κ -(λ', λ')-compact.

(g) Every κ -(λ, μ)-compact logic generated by λ' cardinality quantifiers is (λ', λ') -compact.

Remark 1.4. The equivalence of conditions (a) and (e) above, for $\kappa \geq \sup\{\lambda, \lambda'\}$, has been announced in [L4].

For $\kappa = \lambda = \mu$ and $\lambda' = \mu'$ both regular cardinals, the equivalence of Conditions (e), (f), (g) above has been announced in [L3, p. 80].

Lemma 1.5. *Suppose that $\lambda \geq \mu$ and $\lambda' \geq \mu'$ are infinite cardinals, and κ is any cardinal. Suppose that $(f_\beta)_{\beta \in \kappa}$ is a given set of functions from $S_\mu(\lambda)$ to $S_{\mu'}(\lambda')$.*

Then the following are equivalent.

(a) *Whenever D is an ultrafilter over $S_\mu(\lambda)$ and D covers λ , then there exists some $\beta \in \kappa$ such that $f_\beta(D)$ covers λ' .*

(b) *For every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \subseteq \bigcup_{\beta \in F} f_\beta^{-1}[\{g(\beta)\}]$.*

Proof. We show that the negation of (a) is equivalent to the negation of (b).

Indeed, (a) is false if and only if there exists an ultrafilter D over $S_\mu(\lambda)$ which covers λ and such that, for every $\beta \in \kappa$, $f_\beta(D)$ does not cover λ' .

This means that D covers λ and, for every $\beta \in \kappa$, there exists some $g(\beta) \in \lambda'$ such that $[\{g(\beta)\}] = \{s' \in S_{\mu'}(\lambda') \mid g(\beta) \in s'\} \notin f_\beta(D)$, that is, $f_\beta^{-1}[\{g(\beta)\}] \notin D$, that is, $f_\beta^{-1}[\overline{\{g(\beta)\}}] \in D$, since D is required to be an ultrafilter. Here, $\overline{[\{g(\beta)\}]}$ denotes the complement of $[\{g(\beta)\}]$ in $S_{\mu'}(\lambda')$.

Thus, there exists some ultrafilter D which makes (a) false if and only if there exists some function $g : \kappa \rightarrow \lambda'$ such that the set $\{f_\beta^{-1}[\overline{\{g(\beta)\}}] \mid \beta \in \kappa\} \cup \{[\{\alpha\}] \mid \alpha \in \lambda\}$ has the finite intersection property.

Equivalently, there exists some function $g : \kappa \rightarrow \lambda'$ such that, for every finite $F \subseteq \kappa$ and every finite $G \subseteq \lambda$, $\bigcap_{\beta \in F} f_\beta^{-1}[\overline{\{g(\beta)\}}] \cap \bigcap_{\alpha \in G} [\{\alpha\}] \neq \emptyset$.

Since $\bigcap_{\alpha \in G} [\{\alpha\}] = [G]$, the negation of the above statement is: for every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \cap \bigcap_{\beta \in F} f_\beta^{-1}[\overline{\{g(\beta)\}}] = \emptyset$.

This is clearly equivalent to (b). $\square$

Proof of Theorem 1.3. (a) $\Leftrightarrow$ (b) is immediate from Lemma 1.5.

(b) $\Rightarrow$ (c) For $\alpha \in \lambda'$ and $\beta \in \kappa$, define $C_{\alpha, \beta} = f_\beta^{-1}[\{\alpha\}]$.

(c) $\Rightarrow$ (b) For $\beta \in \kappa$ define $f_\beta : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ by $f_\beta(x) = \{\alpha \in \lambda' \mid x \in C_{\alpha, \beta}\}$.

(c) $\Leftrightarrow$ (c') is immediate by taking complements in $S_\mu(\lambda)$.

(e) $\Rightarrow$ (d) is trivial.

(d) $\Rightarrow$ (a). Let $\mathfrak{S}^+$ be an expansion of $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ witnessing (d).

Without loss of generality, we can assume that $\mathfrak{S}^+$ has Skolem functions (see [CK, Section 3.3]). Indeed, since $\kappa \geq \sup\{\lambda, \lambda'\}$, adding Skolem functions to $\mathfrak{S}^+$ involves adding at most κ new symbols.

Consider the set of all functions $f : U \rightarrow U'$ which are definable in $\mathfrak{S}^+$. Enumerate them as $(f_\beta)_{\beta \in \kappa}$. We are going to show that these functions witness (a).

Indeed, let D be an ultrafilter over $S_\mu(\lambda)$ which covers λ . Consider the D -class Id_D of the identity function on $U = S_\mu(\lambda)$. Since D covers λ , then in the model $\mathfrak{C} = \prod_D \mathfrak{S}^+$ we have that $d(\{\alpha\}) \subseteq Id_D$ for every $\alpha \in \lambda$, where d denotes the elementary embedding. Trivially, $\mathfrak{C} \models U(Id_D)$. Let $\mathfrak{B}$ be the Skolem hull of Id_D in $\mathfrak{C}$. Since $\mathfrak{S}^+$ has Skolem functions, $\mathfrak{B} \preceq \mathfrak{C}$ [CK, Proposition 3.3.2]; in particular, $\mathfrak{B} \equiv \mathfrak{C}$. By Łoś Theorem, $\mathfrak{C} \equiv \mathfrak{S}^+$. By transitivity, $\mathfrak{B} \equiv \mathfrak{S}^+$.

Since $Id_D \in B$, then $\mathfrak{B}$ is (λ, μ) -regular, by what we have proved. Since $\mathfrak{S}^+$ witnesses (d), then $\mathfrak{B}$ has an element x'_D such that $\mathfrak{B} \models U'(x'_D)$ and $\mathfrak{B} \models \{\alpha'\} \subseteq x'_D$ for every $\alpha' \in \lambda'$.

Since $\mathfrak{B}$ is the Skolem hull of Id_D in $\mathfrak{C}$, we have $x'_D = f(Id_D)$, for some function $f : S \rightarrow S$ definable in $\mathfrak{S}^+$, where S is the base set of $\mathfrak{S}^+$.

Since f is definable in $\mathfrak{S}^+$, then also the following function $f' : U \rightarrow U'$ is definable in $\mathfrak{S}^+$:

$$f'(u) = \begin{cases} f(u) & \text{if } u \in U \text{ and } f(u) \in U' \\ \emptyset & \text{if } u \in U \text{ and } f(u) \notin U' \end{cases}$$

Since $\mathfrak{B} \models U'(x'_D)$, and $\mathfrak{B} \preceq \mathfrak{C}$, then $\{u \in U = S_\mu(\lambda) \mid x'(u) \in U'\} \in D$. Since $x'_D = f(Id_D)$, then $\{u \in U \mid x'(u) = f(Id(u)) = f(u)\} \in D$. Hence, $\{u \in U \mid f(u) \in U'\} \in D$, $\{u \in U \mid f(u) = f'(u)\} \in D$ and $\{u \in U \mid x'(u) = f'(u)\} \in D$. Hence, $x'_D = f'_D$.

Since $f' : U \rightarrow U'$ and f' is definable in $\mathfrak{S}^+$, then $f' = f_\beta$ for some $\beta \in \kappa$. We want to show that $D' = f'(D)$ covers λ' .

Indeed, for $\alpha' \in \lambda'$, $[\{\alpha'\}] = \{u' \in S_{\mu'}(\lambda') \mid \alpha' \in u'\} \in f'(D)$ if and only if $\{u \in S_\mu(\lambda) \mid \alpha' \in f'(u)\} \in D$, and this is true for every $\alpha' \in \lambda'$, since $\mathfrak{B} \models \{\alpha'\} \subseteq x'_D$ and $x'_D = f'_D$.

(a) $\Rightarrow$ (e). Suppose we have functions $(f_\beta)_{\beta \in \kappa}$ as given by 1.3(a).

Expand $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ to a model $\mathfrak{S}^+$ by adding, for each $\beta \in \kappa$, a new function symbol representing f_β (by abuse of notation, in what follows we shall write f_β both for the function itself and for the symbol that represents it).

Suppose that $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is (λ, μ) -regular, that is, $\mathfrak{B}$ has an element x such that $\mathfrak{B} \models U(x)$, and $\mathfrak{B} \models \{\alpha\} \subseteq x$ for every $\alpha \in \lambda$.

For every formula $\phi(z)$ with just one variable z in the language of $\mathfrak{S}^+$ let $E_\phi = \{s \in S_\mu(\lambda) \mid \mathfrak{S}^+ \models \phi(s)\}$. Let $F = \{E_\phi \mid \mathfrak{B} \models \phi(x)\}$. Since the intersection of any two members of F is still in F , and $\emptyset \notin F$, then F can be extended to an ultrafilter D on $S_\mu(\lambda)$.

For every $\alpha \in \lambda$, consider the formula $\phi(z) \equiv \{\alpha\} \subseteq z$. We get $E_\phi = \{s \in S_\mu(\lambda) \mid \mathfrak{S}^+ \models \{\alpha\} \subseteq s\} = [\{\alpha\}]$. On the other side, since $\mathfrak{B} \models \{\alpha\} \subseteq x$, then by the definition of F we have $E_\phi = [\{\alpha\}] \in F \subseteq D$. Thus, D covers λ .

By (a), $f_\beta(D)$ covers λ' , for some $\beta \in \kappa$. This means that $[\{\alpha'\}] = \{s' \in S_{\mu'}(\lambda') \mid \{\alpha'\} \subseteq s'\} \in f_\beta(D)$, for every $\alpha' \in \lambda'$. That is, $\{s \in S_\mu(\lambda) \mid \{\alpha'\} \subseteq f_\beta(s)\} \in D$ for every $\alpha' \in \lambda'$.

For every $\alpha' \in \lambda'$ consider the formula $\psi(z) \equiv \{\alpha'\} \subseteq f_\beta(z)$. By the previous paragraph, $E_\psi \in D$. Notice that $E_{\neg\psi}$ is the complement of E_ψ in $S_\mu(\lambda)$. Since D is proper, and $E_\psi \in D$, then $E_{\neg\psi} \notin D$. Since D extends F , and either $E_\psi \in F$ or $E_{\neg\psi} \in F$, we necessarily have $E_\psi \in F$, that is, $\mathfrak{B} \models \psi(x)$, that is, $\mathfrak{B} \models \{\alpha'\} \subseteq f_\beta(x)$.

Moreover, since $f_\beta : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$, $\mathfrak{B} \models U(x)$ and $\mathfrak{B} \equiv \mathfrak{S}^+$, then $\mathfrak{B} \models U'(f_\beta(x))$.

Thus, we have proved that $\mathfrak{B}$ has an element $y = f_\beta(x)$ such that $\mathfrak{B} \models U'(y)$ and $\mathfrak{B} \models \{\alpha'\} \subseteq y$ for every $\alpha' \in \lambda'$. This means that $\mathfrak{B}$ is (λ', μ') -regular.

The equivalence of (f) and (g) with the other conditions shall be proved elsewhere. $\square$

2. A MULTICARDINAL GENERALIZATION

Let us recall some generalizations of Definitions 1.1 and 1.2, generalizations introduced in the statement of [L3, Theorem 0.20].

Definitions 2.1. Recall from Definition 1.1 that an ultrafilter D over $S_\mu(\lambda)$ covers λ if and only if $[\{\alpha\}] \in D$, for every $\alpha \in \lambda$.

Suppose that κ is any cardinal, and $\lambda, \mu, (\lambda_\beta)_{\beta \in \kappa}, (\mu_\beta)_{\beta \in \kappa}$ are infinite cardinals such that $\lambda \geq \mu$, and $\lambda_\beta \geq \mu_\beta$ for every $\beta \in \kappa$.

The relation $(\lambda, \mu) \Rightarrow \bigvee_{\beta \in \kappa} (\lambda_\beta, \mu_\beta)$ holds if and only if there are κ functions $(f_\beta)_{\beta \in \kappa}$ such that, for every $\beta \in \kappa$, $f_\beta : S_\mu(\lambda) \rightarrow S_{\mu_\beta}(\lambda_\beta)$ and such that whenever D is an ultrafilter over $S_\mu(\lambda)$ which covers λ , then for some $\beta \in \kappa$ it happens that $f_\beta(D)$ covers λ_β .

Notice that in the case when $\lambda_\beta = \lambda'$ and $\mu_\beta = \mu'$ for every $\beta \in \kappa$ then $(\lambda, \mu) \Rightarrow \bigvee_{\beta \in \kappa} (\lambda_\beta, \mu_\beta)$ is the same as $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$.

$\mathfrak{S}(\lambda, \mu; \lambda_\beta, \mu_\beta)_{\beta \in \kappa}$ denotes the model $\langle S_{\mu''}(\lambda''), \subseteq, U, U_\beta, \{\alpha\}_{\alpha \in \lambda'', \beta \in \kappa} \rangle$, where $\lambda'' = \sup\{\lambda, \sup_{\beta \in \kappa} \lambda_\beta\}$, $\mu'' = \sup\{\mu, \sup_{\beta \in \kappa} \mu_\beta\}$, $U(x)$ if and only if $x \in S_\lambda(\mu)$, and, for $\beta \in \kappa$, $U_\beta(y)$ if and only if $y \in S_{\lambda_\beta}(\mu_\beta)$.

If $\mathfrak{S}^+$ is an expansion of $\mathfrak{S}(\lambda, \mu; \lambda_\beta, \mu_\beta)_{\beta \in \kappa}$, and $\mathfrak{B} \equiv \mathfrak{S}^+$, then, for $\beta \in \kappa$, we say that $\mathfrak{B}$ is $(\lambda_\beta, \mu_\beta)$ -regular if and only if there is $b \in B$ such that $\mathfrak{B} \models U_\beta(b)$, and $\mathfrak{B} \models \{\alpha\} \subseteq b$ for every $\alpha \in \lambda_\beta$. The definition of a (λ, μ) -regular extension is as in Definition 1.2

Theorem 2.2. *Suppose that κ is any cardinal, and $\lambda, \mu, (\lambda_\beta)_{\beta \in \kappa}, (\mu_\beta)_{\beta \in \kappa}$ are infinite cardinals such that $\lambda \geq \mu$, and $\lambda_\beta \geq \mu_\beta$ for every $\beta \in \kappa$.*

Then the following conditions are equivalent.

- (a) $(\lambda, \mu) \Rightarrow \bigvee_{\beta \in \kappa} (\lambda_\beta, \mu_\beta)$ holds.
- (b) There are κ functions $(f_\beta)_{\beta \in \kappa}$ such that, for every $\beta \in \kappa$, $f_\beta : S_\mu(\lambda) \rightarrow S_{\mu_\beta}(\lambda_\beta)$ and such that for every function $g \in \prod_{\beta \in \kappa} \lambda_\beta$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \subseteq \bigcup_{\beta \in F} f_\beta^{-1}[\{g(\beta)\}]$.
- (c) There is a family $(C_{\alpha, \beta})_{\alpha \in \lambda_\beta, \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:
 - (i) For every $\beta \in \kappa$ and every $H \subseteq \lambda_\beta$, if $|H| \geq \mu_\beta$ then $\bigcap_{\alpha \in H} C_{\alpha, \beta} = \emptyset$.
 - (ii) For every function $g \in \prod_{\beta \in \kappa} \lambda_\beta$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \subseteq \bigcup_{\beta \in F} C_{g(\beta), \beta}$.
- (c') There is a family $(B_{\alpha, \beta})_{\alpha \in \lambda_\beta, \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:
 - (i) For every $\beta \in \kappa$ and every $H \subseteq \lambda_\beta$, if $|H| \geq \mu_\beta$ then $\bigcup_{\alpha \in H} B_{\alpha, \beta} = S_\mu(\lambda)$.
 - (ii) For every function $g \in \prod_{\beta \in \kappa} \lambda_\beta$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq \lambda$ such that $[G] \cap \bigcap_{\beta \in F} B_{g(\beta), \beta} = \emptyset$.

Suppose in addition that $\kappa \geq \lambda$ and $\kappa \geq \lambda_\beta$ for all $\beta \in \kappa$, and that, for every $\beta_0 \in \kappa$, $|\{\beta \in \kappa \mid \lambda_\beta = \lambda_{\beta_0} \text{ and } \mu_\beta = \mu_{\beta_0}\}| = \kappa$. Then the preceding conditions are also equivalent to:

- (d) $\mathfrak{S}(\lambda, \mu; \lambda_\beta, \mu_\beta)_{\beta \in \kappa}$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{S}^+$ with at most κ new symbols and such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is (λ, μ) -regular, then there is $\beta \in \kappa$ such that $\mathfrak{B}$ is $(\lambda_\beta, \mu_\beta)$ -regular.

Suppose further that $\lambda_\beta = \mu_\beta$ is a regular cardinal for every $\beta \in \kappa$. Then the preceding conditions are also equivalent to:

- (e) Every κ -(λ, μ)-compact logic is κ -($\lambda_\beta, \lambda_\beta$)-compact for some $\beta \in \kappa$.
- (f) Every κ -(λ, μ)-compact logic generated by $\sup_{\beta \in \kappa} \lambda_\beta$ cardinality quantifiers is $(\lambda_\beta, \lambda_\beta)$ -compact for some $\beta \in \kappa$.

Proof. There is no essential difference with the proof of Theorem 1.3. $\square$

Notice that Theorem 1.3. is the particular case of Theorem 2.2 when $\lambda_\beta = \lambda'$ and $\mu_\beta = \mu'$ for every $\beta \in \kappa$.

3. THE “ALMOST” GENERALIZATIONS

Versions of Theorem 1.3 can be given for the “almost” variants of $(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$. In order to state the above remark precisely, we need to introduce some variations on Definitions 1.1 and 1.2 (see [L3, Definition 0.14]).

Definition 3.1. We say that an ultrafilter D over $S_\mu(\lambda)$ *almost covers* λ if and only if $|\{\alpha \in \lambda \mid [\alpha] \in D\}| = \lambda$.

The relation $\text{alm}(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ holds if and only if there are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that whenever D is an ultrafilter over $S_\mu(\lambda)$, and D almost covers λ , then for some $\beta \in \kappa$ it happens that $f_\beta(D)$ covers λ' .

The relations $\text{alm}(\lambda, \mu) \xrightarrow{\kappa} \text{alm}(\lambda', \mu')$ and $(\lambda, \mu) \xrightarrow{\kappa} \text{alm}(\lambda', \mu')$ are defined similarly.

Notice that $(\lambda, \mu) \xrightarrow{1} \text{alm}(\lambda, \mu)$ trivially. Moreover, if ν is a regular cardinal, then $\text{alm}(\nu, \nu) \xrightarrow{1} (\nu, \nu)$, as witnessed by $f : S_\nu(\nu) \rightarrow S_\nu(\nu)$ defined by $f(x) = \sup x$ (cf. also [L3, Lemma 0.16(ii)]). Thus, in the next results, if either $\nu = \lambda = \mu$, or $\nu' = \lambda' = \mu'$, or both, regular cardinals, then $\text{alm}(\nu, \nu)$ and (ν, ν) can be used interchangeably. A similar remark applies to almost (ν, ν) -regularity and (ν, ν) -regularity, as defined below.

Recall the definition of the model $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ from Definition 1.2.

If $\mathfrak{S}^+$ is an expansion of $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$, and $\mathfrak{B} \equiv \mathfrak{S}^+$, we say that $\mathfrak{B}$ is *almost (λ, μ) -regular* if and only if there is $b \in B$ such that $\mathfrak{B} \models U(b)$, and $|\{\alpha \in \lambda \mid \mathfrak{B} \models \{\alpha\} \subseteq b\}| = \lambda$. The notion of *almost (λ', μ') -regularity* is defined similarly.

Theorem 3.2. *Suppose that $\lambda \geq \mu$, $\lambda' \geq \mu'$ are infinite cardinals, and κ is any cardinal.*

Then the following conditions are equivalent.

- (a) $(\lambda, \mu) \xrightarrow{\kappa} \text{alm}(\lambda', \mu')$ holds.
- (b) *There are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that for every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq \lambda$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \subseteq \bigcup_{\beta \in F, \beta^* \in H_\beta} f_\beta^{-1}[\{g(\beta^*)\}]$.*
- (c) *There is a family $(C_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:*

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcap_{\alpha \in H} C_{\alpha, \beta} = \emptyset$.

(ii) For every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq \lambda$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \subseteq \bigcup_{\beta \in F, \beta^* \in H_\beta} C_{g(\beta^*), \beta}$.

(c') There is a family $(B_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcup_{\alpha \in H} B_{\alpha, \beta} = S_\mu(\lambda)$.

(ii) For every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq \lambda$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \cap \bigcap_{\beta \in F, \beta^* \in H_\beta} B_{g(\beta^*), \beta} = \emptyset$.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, then the preceding conditions are also equivalent to:

(d) $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{S}^+$ with at most κ new symbols such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is (λ, μ) -regular, then $\mathfrak{B}$ is almost (λ', μ') -regular.

Theorem 3.3. Suppose that $\lambda \geq \mu$, $\lambda' \geq \mu'$ are infinite cardinals, and κ is any cardinal.

Then the following conditions are equivalent.

(a) $\text{alm}(\lambda, \mu) \xrightarrow{\kappa} (\lambda', \mu')$ holds.

(b) There are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that for every $T \subseteq \lambda$ with $|T| = \lambda$, and for every function $g : \kappa \rightarrow \lambda'$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq T$ such that $[G] \subseteq \bigcup_{\beta \in F} f_\beta^{-1}[\{g(\beta)\}]$.

(c) There is a family $(C_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcap_{\alpha \in H} C_{\alpha, \beta} = \emptyset$.

(ii) For every function $g : \kappa \rightarrow \lambda'$ and for every $T \subseteq \lambda$ with $|T| = \lambda$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq T$ such that $[G] \subseteq \bigcup_{\beta \in F} C_{g(\beta), \beta}$.

(c') There is a family $(B_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcup_{\alpha \in H} B_{\alpha, \beta} = S_\mu(\lambda)$.

(ii) For every function $g : \kappa \rightarrow \lambda'$ and for every $T \subseteq \lambda$ with $|T| = \lambda$ there exist finite sets $F \subseteq \kappa$ and $G \subseteq T$ such that $[G] \cap \bigcap_{\beta \in F} B_{g(\beta), \beta} = \emptyset$.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, then the preceding conditions are also equivalent to:

(d) $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{S}^+$ with at most κ new symbols such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is almost (λ, μ) -regular, then $\mathfrak{B}$ is (λ', μ') -regular.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, and $\lambda' = \mu'$ is a regular cardinal then the preceding conditions are also equivalent to:

(f) Every almost κ -(λ, μ)-compact logic is κ -(λ', λ')-compact.

(g) Every almost κ -(λ, μ)-compact logic generated by λ' cardinality quantifiers is (λ', λ') -compact.

Theorem 3.4. Suppose that $\lambda \geq \mu$, $\lambda' \geq \mu'$ are infinite cardinals, and κ is any cardinal.

Then the following conditions are equivalent.

(a) $\text{alm}(\lambda, \mu) \xrightarrow{\kappa} \text{alm}(\lambda', \mu')$ holds.

(b) There are κ functions $(f_\beta)_{\beta \in \kappa} : S_\mu(\lambda) \rightarrow S_{\mu'}(\lambda')$ such that for every $T \subseteq \lambda$ with $|T| = \lambda$ and for every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq T$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \subseteq \bigcup_{\beta \in F, \beta^* \in H_\beta} f_\beta^{-1}[\{g(\beta^*)\}]$.

(c) There is a family $(C_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcap_{\alpha \in H} C_{\alpha, \beta} = \emptyset$.

(ii) For every $T \subseteq \lambda$ with $|T| = \lambda$ and for every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq T$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \subseteq \bigcup_{\beta \in F, \beta^* \in H_\beta} C_{g(\beta^*), \beta}$.

(c') There is a family $(B_{\alpha, \beta})_{\alpha \in \lambda', \beta \in \kappa}$ of subsets of $S_\mu(\lambda)$ such that:

(i) For every $\beta \in \kappa$ and every $H \subseteq \lambda'$, if $|H| \geq \mu'$ then $\bigcup_{\alpha \in H} B_{\alpha, \beta} = S_\mu(\lambda)$.

(ii) For every $T \subseteq \lambda$ with $|T| = \lambda$ and for every function $g : \kappa \rightarrow S_{\lambda'}(\lambda')$ there exist a finite set $G \subseteq T$, a finite set $F \subseteq \kappa$, and, for $\beta \in F$, finite sets $H_\beta \subseteq \lambda' \setminus g(\beta)$ such that $[G] \cap \bigcap_{\beta \in F, \beta^* \in H_\beta} B_{g(\beta^*), \beta} = \emptyset$.

If in addition $\kappa \geq \sup\{\lambda, \lambda'\}$, then the preceding conditions are also equivalent to:

(d) $\mathfrak{S}(\lambda, \mu; \lambda', \mu')$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{S}^+$ with at most κ new symbols such that whenever $\mathfrak{B} \equiv \mathfrak{S}^+$ and $\mathfrak{B}$ is almost (λ, μ) -regular, then $\mathfrak{B}$ is almost (λ', μ') -regular.

Proofs. Similar to the proof of Theorem 1.3. $\square$

Remark 3.5. There is a common generalization of Theorems 2.2 and 3.2. There is a simultaneous generalization of Theorems 3.3 and 3.4 along the lines of Theorem 2.2. We leave details to the reader.

In fact, we have a common generalization of all the results presented in this note, including Remark 4.3 below. Details shall be presented elsewhere.

4. TWO PROBLEMS AND A FURTHER GENERALIZATION

Problem 4.1. It is proved in [L6, Theorem 2] that $\text{alm}(\lambda^+, \mu^+) \xRightarrow{\lambda^+} \text{alm}(\lambda, \mu)$ holds.

Is it true that $(\lambda^+, \mu^+) \xRightarrow{\lambda^+} (\lambda, \mu)$ holds?

This is true when $\lambda = \mu$ is a regular cardinal.

Problem 4.2. As proved in [FMS], it is consistent, modulo some large cardinal assumption, that there is a uniform ultrafilter over ω_1 which is not (ω, ω_1) -regular. This is equivalent to the failure of $(\omega_1, \omega_1) \xRightarrow{2^{\omega_1}} (\omega_1, \omega)$.

Is it possible to find a model for the failure of $(\omega_1, \omega_1) \xRightarrow{\omega_1} (\omega_1, \omega)$ by using weaker consistency assumptions?

Which is the exact consistency strength of the failure of $(\omega_1, \omega_1) \xRightarrow{\omega_1} (\omega_1, \omega)$?

More generally, for arbitrary λ , which is the exact consistency strength of the failure of $(\lambda^+, \lambda^+) \xRightarrow{\lambda^+} (\lambda^+, \lambda)$?

Remark 4.3. We can extend the definitions and the results of the present note as follows. If λ and μ are *ordinals*, let $S_\mu(\lambda)$ denote the set of all subsets of λ having order type $< \mu$. See also [BK].

Definitions 1.1, 1.2, 2.1 and 3.1 can be easily generalized to the case when λ and μ are ordinals. In the present situation, the most appropriate definition of “almost covering” appears to be the following: an ultrafilter D over $S_\mu(\lambda)$ almost covers λ if and only if order type of $\{\alpha \in \lambda \mid [\alpha] \in D\} = \lambda$.

The definition of κ -(λ, μ)-compactness for logics, too, can be easily generalized when λ and μ are ordinals, by always taking into account order type, instead of cardinality.

Notice that, in all the definitions and the results here, κ is used just as an index set; the cardinal structure on κ is not used at all. Hence, allowing κ to be an ordinal is no gain in generality.

All the results of the present note, when appropriately formulated, extend to the more general setting when $\lambda, \mu, \lambda', \mu', \lambda_\beta, \mu_\beta$ are ordinals.

Everywhere, cardinality assumptions must be replaced by assumptions about order type. For example, the condition $|H| \geq \mu'$ in Theorem 1.3 (c), (c') has to be replaced by “the order type of H is $\geq \mu'$ ”.

The condition $\kappa \geq \sup\{\lambda, \lambda'\}$ before clause (d) in Theorem 1.3 can be replaced by $\kappa \geq \sup\{|\lambda|, |\lambda'|\}$.

The same applies to the condition before clause (f), and we actually need the requirement that $\lambda' = \mu'$ is a regular *cardinal*.

Similar remarks apply to Theorems 2.2 and 3.3.

Theorems 3.2 and 3.4 hold, too, with slight further modifications.

REFERENCES

- [BF] J. Barwise, S. Feferman (editors), *Model-theoretic logics*. Perspectives in Mathematical Logic. Springer-Verlag, New York, 1985.
- [BK] M. Benda and J. Ketonen, Regularity of ultrafilters, *Israel J. Math.* **17**, 231–240 (1974).
- [CK] C. C. Chang and J. Keisler, *Model Theory*, Amsterdam (1977).
- [CN] W. Comfort, S. Negrepointis, *The Theory of Ultrafilters*, Berlin (1974).
- [FMS] M. Foreman, M. Magidor and S. Shelah, Martin’s Maximum, saturated ideals and non-regular ultrafilters. Part II, *Annals of Mathematics* **127**, 521–545, (1988).
- [KM] A. Kanamori and M. Magidor, The evolution of large cardinal axioms in Set Theory, in: *Higher Set Theory*, edited by G. H. Müller and D. S. Scott, 99–275, Berlin (1978).
- [L1] P. Lipparini, *About some generalizations of (λ, μ) -compactness*, Proceedings of the 5th Easter conference on model theory (Wendisch Rietz, 1985), Seminarber., Humboldt-Univ. Berlin, *Sekt. Math.* **93**, 139–141 (1987). Available also at the author’s web page.
- [L2] P. Lipparini, *The compactness spectrum of abstract logics, large cardinals and combinatorial principles*, *Boll. Unione Matematica Italiana ser. VII*, **4-B** 875–903 (1990).
- [L3] P. Lipparini, *Ultrafilter translations, I: (λ, λ) -compactness of logics with a cardinality quantifier*, *Arch. Math. Logic* **35**, 63–87 (1996).
- [L4] P. Lipparini, *Regular ultrafilters and $[\lambda, \lambda]$ -compact products of topological spaces* (abstract), *Bull. Symbolic Logic* **5** (1999), 121.
- [L5] P. Lipparini, *Combinatorial and model-theoretical principles related to regularity of ultrafilters and compactness of topological spaces. I*, arXiv:0803.3498; *II*:0804.1445; *III*:0804.3737; *IV*:0805.1548 (2008).
- [L6] P. Lipparini, *Every (λ^+, κ^+) -regular ultrafilter is (λ, κ) -regular*, *Proc. Amer. Math. Soc.* **128** (1999), 605–609.

DIPARTIMENTO DI MATEMATICA, VIALE DELLA RICERCA SCIENTIFICA, II
UNIVERSITOMA DI ROMA (TOR VERGATA), I-00133 ROME ITALY

URL: <http://www.mat.uniroma2.it/~lipparin>